

Inclusion criteria:
Any of the following:

- Volume equates to saturating a heavy absorbance pad/tampon more frequently than every 2 hours
- Excessive menstrual flow in duration of ≥ 7 days
- Bleeding is causing symptomatic anemia or lifestyle disturbance
- Bleeding unlikely to be due to ongoing contraception

Exclusion criteria:

- Hormonal bleeding with contraception
- Pregnancy
- Trauma
- Sexual assault
- Genital injury
- Known bleeding disorders

Urgent Care/Express Care Considerations:


- Recommended labs and imaging are not available at CM UC/EC locations

Bleeding Disorder Risk Assessment Checklist*:
Has the pt experienced any of the following:

1. Duration of menses ≥ 7 days?
2. Sensation of flooding or gushing during menses?
3. Need to change menstrual product more frequently than every 2 hours?
4. Impairment of daily activities due to heavy bleeding?
5. Prolonged bleeding with first menses (≥ 7 days)?
6. History of excessive bleeding with tooth extraction or surgery?
7. If history of pregnancy, experienced excessive bleeding following delivery or miscarriage?
8. History of blood transfusion or medication for anemia due to menstrual bleeding?
9. Family history of a diagnosed bleeding disorder?

If "Yes" to ANY questions, proceed to High Risk for Bleeding Disorder Assessment.

**Adapted from The American College of Obstetrics and Gynecology screening tool*



This clinical pathway is meant as a guide for the healthcare provider, does not establish a standard of care, and is not a substitute for medical judgment which should be applied based upon the individual circumstances and clinical condition of the patient. Copyright © The Children's Mercy Hospital 2023. All rights reserved.