




Screening Criteria: Pts. <u>Not</u> Toilet Trained	
Female Risk Factors • Temp $\geq 39^{\circ}\text{C}$ • Fever ≥ 2 days • No alternative source • ≤ 12 months of age If 2 factors present: ▸ Consider screening If 3 or more factors present ▸ Recommend screening	Male Risk Factors • Temp $\geq 39^{\circ}\text{C}$ • Fever ≥ 2 days • No alternative source • ≤ 6 months of age If 1 factor present: ▸ If uncircumcised, consider screening If 2 factors present: ▸ If circumcised, consider screening ▸ If uncircumcised, recommend screening If 3 or more factors present ▸ If circumcised, recommend screening
Screening Criteria: Pts <u>that are</u> fully Toilet Trained with <u>ANY</u> of the following:	
• Referable urinary tract symptoms ▪ Urinary frequency or urgency ▪ Dysuria ▪ Nausea/vomiting ▪ Urinary incontinence ▪ Abdominal/flank pain ▪ Hematuria • Fever ≥ 2 days without a source in pts with prior UTI history • Fever ≥ 5 days without a source	
Additional Recommendations for Adolescents	
• Collect 'dirty' urine (not a clean catch specimen) for Gonococcus (GC) / Chlamydia (Chl) screening ◦ If GC/Chl positive: Recommend Syphilis Screen • HSV testing: Culture visible lesions, or cervical culture as indicated • Recommend HIV testing • For females: Consider pregnancy testing	
Suspected UTI or Pyelonephritis	
• Clinical signs and symptoms AND • Urinalysis positive for: ◦ WBC ≥ 5 per HPF, OR ◦ Positive leukocyte esterase, OR ◦ Positive nitrite (link to evidence)	

Antibiogram link

Renal Imaging for UTI/Pyelo Algorithm



QR Code for mobile access